

Effects of Exercise on Health-Related Outcomes in Those with Cancer

What can exercise do?

- **Prevention of 7 common cancers***

Dose: 2018 Physical Activity Guidelines for Americans: 150-300 min/week moderate or 75-150 min/week vigorous aerobic exercise









- **Survival of 3 common cancers****

Dose: Exact dose of physical activity needed to reduce cancer-specific or all-cause mortality is not yet known; Overall more activity appears to lead to better risk reduction

*bladder, breast, colon, endometrial, esophageal, kidney and stomach cancers

**breast, colon and prostate cancers

Overall, avoid inactivity, and to improve general health, aim to achieve the current physical activity guidelines for health (150 min/week aerobic exercise and 2x/week strength training).

Outcome	Aerobic Only	Resistance Only	Combination (Aerobic + Resistance)
Strong Evidence	Dose	Dose	Dose
 Cancer-related fatigue	3x/week for 30 min per session of moderate intensity	2x/week of 2 sets of 12-15 reps for major muscle groups at moderate intensity	3x/week for 30 min per session of moderate aerobic exercise, plus 2x/week of resistance training 2 sets of 12-15 reps for major muscle groups at moderate intensity
 Health-related quality of life	2-3x/week for 30-60 min per session of moderate to vigorous	2x/week of 2 sets of 8-15 reps for major muscle groups at a moderate to vigorous intensity	2-3x/week for 20-30 min per session of moderate aerobic exercise plus 2x/week of resistance training 2 sets of 8-15 reps for major muscle groups at moderate to vigorous intensity
 Physical Function	3x/week for 30-60 min per session of moderate to vigorous	2-3x/week of 2 sets of 8-12 reps for major muscle groups at moderate to vigorous intensity	3x/week for 20-40 min per session of moderate to vigorous aerobic exercise, plus 2-3x/week of resistance training 2 sets of 8-12 reps for major muscle group at moderate to vigorous intensity
 Anxiety	3x/week for 30-60 min per session of moderate to vigorous	Insufficient evidence	2-3x/week for 20-40 min of moderate to vigorous aerobic exercise plus 2x/week of resistance training of 2 sets, 8-12 reps for major muscle groups at moderate to vigorous intensity
 Depression	3x/week for 30-60 min per session of moderate to vigorous	Insufficient evidence	2-3x/week for 20-40 min of moderate to vigorous aerobic exercise plus 2x/week of resistance training of 2 sets, 8-12 reps for major muscle groups at moderate to vigorous intensity
 Lymphedema	Insufficient evidence	2-3x/week of progressive, supervised, program for major muscle groups does not exacerbate lymphedema	Insufficient evidence
Moderate Evidence			
 Bone health	Insufficient evidence	2-3x/week of moderate to vigorous resistance training plus high impact training (sufficient to generate ground reaction force of 3-4 time body weight) for at least 12 months	Insufficient evidence
 Sleep	3-4x/week for 30-40 min per session of moderate intensity	Insufficient evidence	Insufficient evidence

Citation: bit.ly/cancer_exercise_guidelines

Moderate intensity (40%-59% heart rate reserve or VO₂R) to vigorous intensity (60%-89% heart rate reserve or VO₂R) is recommended.

Exercise is Medicine®

AMERICAN COLLEGE of SPORTS MEDICINE